

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 89104464.6


 (51) Int. Cl.4: **C07D 471/04 , A61K 31/435 ,**  
**/(C07D471/04,235:00,221:00)**

(22) Date of filing: 14.03.89

<p>Amended claims in accordance with Rule 86 (2) EPC.</p> <p>(30) Priority: 14.03.88 US 167671 06.03.89 US 317871</p> <p>(43) Date of publication of application: 06.12.89 Bulletin 89/49</p> <p>(84) Designated Contracting States: <b>AT BE CH DE ES FR GB GR IT LI LU NL SE</b></p> <p>(71) Applicant: G.D. Searle &amp; Co. P.O. Box 5110 Chicago Illinois 60680(US)</p>	<p>(72) Inventor: Khanna, Ish K. 8922 Bronx Avenue Skokie, Il. 60077(US) Inventor: Nosal, Roger 661 Indian Spring Lane Buffalo Grove, Il. 60089(US) Inventor: Weier, Richard Mathias 240 Hickory Court Lake Bluff, Il. 60044(US)</p> <p>(74) Representative: Beil, Hans Chr., Dr. et al Beil, Wolff und Beil, Rechtsanwälte Adelonstrasse 58 Postfach 80 01 40 D-6230 Frankfurt am Main 80(DE)</p>
--	--

(54) 5-Substitued (4,5-c)imidazopyridines.

(57) This invention relates to novel substituted imidazopyridine dervatives having the following formula



or a pharmaceutically acceptable acid addition salt: wherein

R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; heterocyclical-  
 kyl having 4 to 8 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; heteroaromatic having 5 or 6 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen; R<sub>1</sub> and R<sub>2</sub> cannot both be hydrogen

Y is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the

**EP 0 344 414 A1**

64

phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxyalkyl wherein the alkyl groups are each 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl group are each 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms; and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.

n is an integer of 1 to 5.

R<sub>3</sub> is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms.

useful in the treatment of diseases or disorders mediated by platelet-activating factor. This invention relates to pharmaceutical compositions of such substituted imidazopyridines.

## 5-Substituted [4,5-c] Imidazopyridines

This application is a continuation-in-part of serial number 07/167,671, filed March 14, 1988, now pending.

FIELD OF THE INVENTION

This invention is in the field of mammalian therapeutics and relates to compounds for treatment of mammalian diseases such as inflammation, cardiovascular disorders, asthma and other diseases. Of particular interest is a class of 5-substituted [4,5-c] imidazopyridines useful for treatment of cardiovascular and immuno-inflammatory related disorders mediated by platelet activating factor (PAF).


BACKGROUND OF THE INVENTION

Platelet-activating factor (PAF) has been associated with various biological activities and pathways, thus making it an important mediator responsible for a variety of physiological processes including, but not limited to, activation and aggregation of platelets, smooth muscle contraction, pathogenesis of immune complex deposition, inflammation, and respiratory, cardiovascular and intravascular alterations. These physiological processes are associated with a large group of diseases, such as, for example, cardiovascular disorders, asthma, lung edema, endotoxin shock, adult respiratory distress syndrome and inflammatory diseases.

United States Patent 4,804,658 discloses a class of imidazopyridine derivatives useful in the treatment of diseases or disorders mediated by platelet-activating factor. The present invention is distinct from this disclosure in that in the present invention the benzamide moiety is attached to the nitrogen (position 5) which makes up the six membered ring of the imidazopyridine ring system as opposed to the disclosure wherein the benzamide moiety is attached to one of the nitrogens which makes up the five membered ring of the imidazopyridine ring system.

Summary of the Invention

The present invention relates to a novel class of compounds represented by the formula



or a pharmaceutically acceptable acid addition salt thereof; wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; heterocycloalkyl having 4 to 8 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; heteroaromatic having 5 or 6 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of

the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms substituted straight or branched chain alkyl which can be substituted one or more by halogen; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxyalkyl wherein the alkyl groups are each 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl group are each 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms; and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.


$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

The invention further relates to pharmaceutical compositions comprising a compound of formula I. Such compounds and compositions have potent and specific PAF antagonistic activities and are thereby useful in the treatment of various diseases or disorders mediated by PAF, for example inflammation, cardiovascular disorders, asthma, lung edema, and adult respiratory distress syndrome.

A preferred embodiment of the present invention are compounds of the formula



or a pharmaceutically acceptable acid addition salt thereof; wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen


$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen;

$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6 or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

A further embodiment of the present invention are compounds of the formula



or a pharmaceutically acceptable acid addition salt thereof, wherein

$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by group independently selected from alkyl of 1 to 6 carbon atoms or halogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms;

$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

As used herein the term "alkyl of 1 to 15 carbon atoms": refers to straight chain or branched chain hydrocarbon groups having from one to fifteen carbon atoms. Illustrative of such alkyl groups are methyl, ethyl, propyl isopropyl, butyl, isobutyl, pentyl, neopentyl, hexyl, isohexyl, octyl, decyl and the like.

As used herein the term "cycloalkyl having 3 to 8 carbon atoms" included cycloalkyl groups having from three to eight carbons. Illustrative of such cycloalkyl groups are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and the like.

As used herein the term halogen includes fluoro, chloro and bromo.

As used herein the term "alkenyl having 2 to 15 carbon atoms" refers to straight or branched unsaturated hydrocarbon groups having from 2 to 15 carbon atoms. Illustrative of such alkenyl groups are 2-propenyl, hexenyl, octenyl, decenyl and the like.

As used herein the term "alkoxy wherein the alkyl is 1 to 6 carbon atoms" refers to straight or branched chain ethers. Illustrative of such groups are methoxy, ethoxy, propoxy, butoxy, isopropoxy and the like.

The term "hydroxyalkyl" refers to straight or branched alkyl group having one to six atoms any one of which may be substituted with one or more hydroxyl group.

The term "thioalkyl" refers to straight or branched thio-containing radicals, respectively having alkyl portions of one to six attached.

The term "mercaptoalkyl" refers to a terminal mercapto group attached to an alkyl portion of one to six carbon atoms which can be straight or branched.


The term "heterocyclicalkyl" refers to a cyclic radicals having 5 to 8 carbon atoms wherein one or more of the carbons is replaced by nitrogen, sulfur or oxygen.

The term heteroaromatic refers to cyclic radicals having 5 or 6 ring carbon atoms which can be optionally substituted one or more times by alkyl of 1 to 6 carbon atoms with the understanding that the 5 membered carbon atom ring is replaced one or more times by nitrogen, sulfur or oxygen and when more than one hetero atom exists in the 5-membered ring one hetero atom must be nitrogen; the six membered carbon atom ring is replaced one or more times by nitrogen.


Included within the embodiments of the present invention are the tautomeric forms of the described compounds, isomeric forms including geometric isomers, enantiomers and diastereoisomers, and the pharmaceutically acceptable salts thereof.

The term "pharmaceutically acceptable acid addition salt" refers to a salt prepared by contacting a compound of formula (I) with an acid whose anion is generally considered suitable for human consumption. Examples of pharmacologically acceptable acid addition salts include the hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate, acetate, propionate, lactate, maleate, malate, succinate, and tartrate salts. All of these salts may be prepared by conventional means by reacting, for example, the appropriate acid with the corresponding compound of Formula I.

The compounds of formula (I) may be prepared in accordance with the following procedures.  
Imidazopyridine which is represented by the following formula



wherein  $R_3$  and  $R_4$  are defined as before is reacted with a haloalkylbenzamide which is represented by the following formula




wherein  $R_1$  and  $R_2$  and  $n$  are defined as before and  $X$  is chloro, bromo, or methanesulfonyloxy to give the compounds of formula I. It is understood that the haloalkylbenzamide can also be substituted by halogen, alkyl of 1 to 6 carbon atoms; alkoxy wherein the alkyl is 1 to 6 carbon atoms; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxy alkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl is 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.

Preferred reaction conditions for the above-identified procedure include heating overnight at 70-90 °C a solution of haloalkylbenzamide and imidazopyridine in a solvent such as dimethylacetamide (approximately 0.1M in each). After heating overnight the reaction solvent is removed in vacuo and the residue diluted with water and basified with ammonium hydroxide. The aqueous solution is extracted with chloroform and the combined organic extracts are backwashed with saturated aqueous sodium chloride solution. The organic solution is dried over sodium sulfate or magnesium sulfate, the drying agent filtered and the filtrate concentrated in vacuo to give the crude product. Purification is effected by chromatography on silica gel using mixtures of chloroform, ethanol and ammonium hydroxide.

A preferred work up for the above-described procedure is to cool the reaction solution which had been heated overnight to room temperature and remove the solvent under reduced pressure at <45 °C. The residue obtained is triturated with excess of dry ether and filtered. The crude product is purified by chromatography.

#### Preparation of Intermediates


## Scheme A



The imidazo [4,5-c]pyridine wherein  $R_3$  is 4-methyl is prepared according to the scheme above starting with the imidazopyridine of Formula II. Position 1 of this compound is protected by reaction with a 2-(trialkylsilyl) ethoxy methyl chloride and a base such as sodium hydride or potassium hydride in a polar aprotic solvent such as dimethylformamide. This reaction is carried out at room temperature. A specific example of such a protecting reagent is 2-(trimethylsilyl)ethoxymethyl chloride. The protected imidazopyridine is reacted with phenyl chloroformate and methylmagnesium bromide in an ether solvent such as tetrahydrofuran at about  $-20^\circ\text{C}$ . The methylated product bearing phenoxycarbonyl at position 5 is treated with a base, such as alcoholic sodium hydroxide, at reflux for 24 hr. The product is oxidized with, for example, chloranil, and the 2-(trimethylsilyl)ethoxymethyl group is removed by treatment with a suitable acid. An example of such an acid would be trifluoroacetic acid. Preparation of the unsubstituted imidazo [4,5-c] pyridine is described in U. S. Patent No. 4,804,658.

The haloalkyl benzamides are prepared according to the following reaction scheme

## Scheme B



wherein  $R_1$  and  $R_2$  are defined as before; Z is  $\text{CH}_2\text{Br}$  or H; X is fluoro, OMe or methyl

Thus according to the above scheme the acid chlorides were prepared from the corresponding carboxylic acids by refluxing in thionyl chloride (2 molar excess) for two hours. Excess thionyl chloride was


removed by azeotrope with toluene. The residual acid chloride was dissolved in THF and cooled to  $-10^{\circ}\text{C}$ . A solution of two molar equivalents of the secondary amine in the THF was added dropwise with stirring. When addition was completed, the reaction was allowed to warm to room temperature and stirred for 1-2 hours. The reaction was quenched with 1N HCL, diluted with  $\text{H}_2\text{O}$  and extracted three times with ethyl acetate. The combined organic layers were washed with saturated aqueous sodium bicarbonate solution, with water and with saturated aqueous sodium chloride and dried over sodium sulfate. The drying agent was filtered and the filtrate concentrated in vacuo to give a crude product that was chromatographed on silica gel using mixtures of ethyl acetate and hexane to give the purified amide.

When  $\text{Z} = \text{CH}_2\text{Br}$  and  $\text{X} = \text{H}$ , the above description is sufficient for the preparation of the compounds of Formula III. When  $\text{Z} = \text{CH}_3$ , and  $\text{X} = \text{OMe}$  or  $\text{F}$ , or when  $\text{Z} = \text{H}$  and  $\text{X} = \text{CH}_3$  then compound of Formula VI must be treated with a halogenating agent such as N-bromo succinimide.

A stirred mixture of the purified amide and NBS (1:1 molar ratio) in carbon tetrachloride was irradiated with a sun lamp for 1-3 hours. A white precipitate was filtered and washed with a minimum amount of  $\text{CHCl}_3$ . The filtrate was washed with water and the aqueous layer, after basification with ammonium hydroxide, was extracted three times with chloroform. All organic layers were combined, washed three times with saturated aqueous sodium chloride solution and dried over sodium sulfate.

The drying agent was filtered and the filtrate concentrated in vacuo to give a crude product that was chromatographed on silica gel using mixtures of ethyl acetate and hexane to give the purified bromomethyl compound.

Scheme C



The benzamides wherein  $n = 2$  or  $3$  can be prepared according to the scheme above starting with the appropriate hydroxyalkyl bromobenzene. The hydroxyl group was protected as a trialkylsilyl ether by reaction with a trialkylsilyl chloride and imidazole in a suitable solvent such as dimethylformamide. An example of such a protecting group would be the *t*-butyldimethylsilyl ether. The crude silyl ether was purified by chromatography on silica gel using mixtures of ethyl acetate and hexane. The aryl bromide was converted to the carboxamide according to the procedure of Schoenberg et al. [J. Org. Chem., 39, 3327- (1974)]. Thus, the aryl bromide was reacted with carbon monoxide in the secondary amine as solvent using bistrisphenylphosphine palladium(II) dibromide as catalyst at about  $100^{\circ}\text{C}$  for 8-26 hr. in a pressure vessel. The reaction vessel was vented, the reaction mixture triturated with ethyl ether and the washings filtered. The filtrate was washed with 10% aqueous HCl, water and brine. After drying over a suitable drying agent, such as magnesium sulfate, and filtering, the filtrate was concentrated and the residue chromatographed on



silica gel using mixtures of ethyl acetate and hexane as eluent to give pure product. The silyl ether was removed by reaction with tetra-n-butylammonium fluoride and the alcohol was converted to a sulfonate ester by reaction with an alkyl or arylsulfonyl chloride. An example of such a sulfonate would be the methanesulfonate.

5 The secondary amines may be prepared by any number of methods known to those skilled in the art. See references

- Emerson, W. S. Org. Reactions 4, 174 1948)


- J. B. Cambell, L. B. Lavaginino in "Catalysis in Organic Synthesis" (Jones W. H., ed.) p. 43, Academic Press, New York, 1980.

10

#### Preparation of 4-Methyl-7-methoxy-imidazopyridine

15

20



25

The above compound can be prepared by the following

30

35


40

45

50


55

Scheme D




Thus the addition product 2 which is isolated on reacting 1 with phenyl chloroformate and methylmagnesium bromide is treated with osmium tetroxide in aqueous acetone containing N-methylmorpholine-N-oxide at room temperature for 24 hours to give the diol 3 and the hydroxyketone 4. The hydroxyketone 4 is acetylated (acetic anhydride, DMAP methylene chloride, room temperature, 24 hrs) and treated with  $\text{CrCl}_2$  in acetone to give the deacetylated product 6. Product 6 is treated with NaH in DMF and then with iodomethane to give the methyl ether 7. Cleavage of carbamate and oxidation gives the N-1 protected 4-methyl-7-methoxyimidazopyridine product 8. Deprotection of product 8 gives the 4-methyl-7-methoxyimidazopyridine.

Preparation of 2-Methoxy-4-bromomethyl-5-bromobenz(N-cyclopentyl,N-2-methylcyclohexyl)amide




The above compound is prepared from 2-methoxy-4-methylbenz(N-cyclopentyl,N-2-methylcyclohexyl)-amide and N-bromo succinimide in carbon tetrachloride by irradiation with a sun lamp for 5 hours.

#### Preparation of 2,6-Dimethoxy-3 bromo-4-bromomethylbenz (N-cyclohexyl,N-cyclopentyl)amide




The above compound is prepared from 2,6-dimethoxy-4-methyl benzoic acid described by I. W. Mathison, R. C. Gueldner, D. M. Carroll, J. Pharma Sci 57 1820, (1968). The substituted benzoic acid is converted to the corresponding amide by first converting said compound to the acid chloride (using thionyl chloride) followed by condensation with N-cyclohexyl,N-cyclopentylamine. Irradiation of 2,6-dimethoxy-4-methyl-benz(N-cyclohexyl, N-cyclopentyl)amide following the procedure described for the preparation 2-methoxy-4-bromomethyl-5-bromobenz(N-cyclopentyl,N-2 methylcyclohexyl)amide gives two products, 2,6-dimethoxy-3-bromo-4-methylbenz-(N-cyclohexyl, N-cyclopentyl)amide and 2,6-dimethoxy-3-bromo-4-bromomethylbenz(N-cyclohexyl,N-cyclopentyl)amide. The product is the predominate product.

## Scheme E



The imidazo[4,5-c]pyridine wherein  $R_3$  is 4-chloro is prepared according to Scheme E starting with the imidazopyridine of Formula II. Position 1 of this compound is protected by reaction with a 2-(trialkylsilyl)-ethoxy methyl chloride and a base such as sodium hydride or potassium hydride in a polar aprotic solvent such as dimethylformamide. The reaction is carried out at room temperature. The protected imidazopyridine is reacted with *m*-chloroperbenzoic acid in methylene chloride at room temperature to give the pyridine N-oxide product. The N-oxide product is heated in  $\text{POCl}_3$  at  $90^\circ\text{C}$  to give 4-chloro-1-chloromethyl imidazopyridine. Treatment of this compound with sodium methoxide in methanol gave the 4-chloro-1-methoxy ethyl imidazopyridine. Reacting this compound with water/acid with heating gave the 4-chloro-imidazo[4,5-c] pyridine.

## Preparation of Alkoxyalkyls



wherein  $R_1$  and  $R_2$  are defined as before; "Hal" is halogen; Z is alkoxy, thioalkyl, mercapto, hydroxy, halo, amino, alkyl and dialkylamino; and  $Z'$  chloro, bromo, methanesulfonyloxy or p-toluenesulfonyloxy.

When Y is substituted with alkoxyalkyl, such substitution may be carried out by methods known to those skilled in the art. Such a method might, for example, employ the substituted benzoic acid 1 (F. Fichter, G. Shetty, *Helv. Chim. Acts*, 20, 563 (1937)) as starting material. This is converted to the amide 2 by first converting acid 1 to the acid chloride by contact with agents such as oxalyl chloride or thionyl chloride and then treating the acid chloride with the desired amine. Amide 2 is converted to halide 3 by treatment with a halogenating agent such as N-bromosuccinimide. Halide 3 is versatile and in addition to serving as an intermediate to alkoxyalkyl compounds, is also an intermediate to alkylthioalkyl, hydroxyalkyl, mercaptoalkyl and alkylaminoalkyl compounds by treatment with the appropriate Z derivative. When halogen is displaced with a metal alkoxide, such as sodium methoxide, the methoxymethyl derivative (4, Z = OMe) is obtained. Conversion of 4 (Z = OMe) to aldehyde 5 (Z = OMe) is effected by controlled reduction with a reducing agent such as diisobutylaluminum hydride, followed by acid hydrolysis. Reduction of aldehyde 5 to alcohol 6 is effected by a second reduction with another reducing agent such as sodium borohydride or lithium tri-t-butoxyaluminum hydride. Alcohol 6 is converted to a derivative suitable for nucleophilic displacement such as 7 where  $Z'$  is a leaving group such as halide or aryl or alkyl sulfonate. Such conversion is effected by treatment of 6 with, for example, p-toluenesulfonyl chloride, methanesulfonyl chloride, or thionyl chloride.

Compounds where Y of formula I is substituted with hydroxy can be made from the corresponding methoxy substituted compounds by treatment with a demethylating reagent such as lithium ethyl mercaptide in a dipolar, aprotic solvent such as dimethylformamide at temperatures ranging from room temperature to 200°.

This invention also relates to a method of treatment for patients (or mammalian animals raised in the dairy, meat, or fur industries or as pets) suffering from disorders or diseases which can be attributed to PAF

as previously described, and more specifically, a method of treatment involving the administration of compound (I) as the active ingredient.

Accordingly, compound (I) can be used among other things to reduce inflammation, to correct respiratory, cardiovascular, and intravascular alterations or disorders, and to regulate the activation or coagulation of platelets, the pathogenesis of immune complex depositions and smooth muscle contractions.

For the treatment of inflammation, cardiovascular disorder, asthma, or other diseases mediated by PAF, compound (I) may be administered orally, topically, parenterally, or by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques.

The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. Therapeutically effective doses of the compounds of the present invention required to prevent or arrest the progress of the medical condition are readily ascertained by one of ordinary skill in the art.

Accordingly, the invention provides a class of novel pharmaceutical compositions comprising one or more compounds of the present invention in association with one or more non-toxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and if desired other active ingredients. The compounds and composition may for example be administered intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit contained in a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. These may with advantage contain an amount of active ingredient from about 1 to 250 mg preferably from about 25 to 150 mg. A suitable daily dose for a mammal may vary widely depending on the condition of the patient and other factors. However, a dose of from about 0.1 to 3000 mg/kg body weight, particularly from about 1 to 100 mg/kg body weight may be appropriate.

The active ingredient may also be administered by injection as a composition wherein, for example, saline, dextrose or water may be used as a suitable carrier. A suitable daily dose is from about 0.1 to 100 mg/kg body weight injected per day in multiple doses depending on the disease being treated. A preferred daily dose would be from about 1 to 30 mg/kg body weight.

The dosage regimen for treating an infectious disease condition with the compounds and/or compositions of this invention is selected in accordance with a variety of factors, including the type, age, weight, sex and medical condition of the patient; the severity of the infection; the route of administration; and the particular compound employed and thus may vary widely.

For therapeutic purposes, the compound of this invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If per os, the compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulphuric acids, gelatin, acacia, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and thus tableted or encapsulated for convenient administration. Alternatively, the compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art. Appropriate dosages, in any given instance, of course depend upon the nature and severity of the condition treated, the route of administration, and the species of mammal involved, including its size and any individual idiosyncrasies.

Representative carriers, diluents and adjuvants include for example, water, lactose, gelatin, starches, magnesium stearate, talc, vegetable oils, gums, polyalkylene glycols, petroleum jelly, etc. The pharmaceutical compositions may be made up in solid form such as granules, powders or suppositories or in a liquid form such as solutions, suspensions or emulsions. The pharmaceutical compositions may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional pharmaceutical adjuvants such as preservatives, stabilizers, wetting agents, emulsifiers, buffers, etc.

Dosage levels of the order from about 1 mg to about 100 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (from about 50 mg to about 5 gs. per patient per day). For example, inflammation is effectively treated and anti-pyretic and analgesic activity manifested by the administration from about 25 to about 75 mg of the compound per kilogram of body weight per day (about 75 mg to about 3.75 gm per patient per day). Preferably, from about 5 mg to about 50 mg per kilogram of body weight per daily dosage produces highly effective results (about 250 mg to about 2.5 gm per patient per day).


The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration of humans may contain from 5 mg to 95 mg of active agent compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to 95 percent of the total composition. Dosage unit forms will generally contain between from about 25 mg to about 500 mg of active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following Examples are intended to further illustrate the present invention and not to limit the invention in spirit or scope. In the Examples, all parts are parts by weight unless otherwise expressly set forth.

#### EXAMPLE 1

5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine




To a stirred solution of imidazopyridine (5.86 g, 49.2 mmol) in DMF (125 ml) under a nitrogen atmosphere was added washed, dried sodium hydride (prepared from 3.54 g of 50% dispersion oil by washing four times with 50-75 ml portions of hexane). After stirring for 1 hr at room temperature, the evolution of hydrogen gas had ceased and the reaction was cooled to  $-10^{\circ}\text{C}$ . N-Methyl-N-cyclohexyl- $\alpha$ -bromo-p-toluyil amide (16.9 g, 54.5 mmol) was added. The reaction was stirred at  $0^{\circ}$  for 45 min. and at room temperature for 3 hrs.

DMF was removed in vacuo and the residue was diluted with  $\text{H}_2\text{O}$  (200 ml) and the resulting solution was saturated with sodium chloride. The aqueous solution was extracted four times with ethyl acetate (100 ml portions) and the combined organic layers were washed three times with saturated aqueous sodium chloride solution (150 ml portions). After drying over sodium sulfate, the organic solution was filtered and concentrated in vacuo to give 13.38 g of crude product as a brown gum. This material was chromatographed on silica gel using ethanol/chloroform/ammonium hydroxide (20/79/1) to give 3.13 g of compound as an orange oil that crystallized on treatment with ethyl acetate. Recrystallization from ethyl acetate yielded 1.06 g.

Analysis Calcd for  $\text{C}_{21}\text{H}_{24}\text{N}_4\text{O}$ .  $1/4\text{H}_2\text{O}$ : C, 71.46; H, 7.00; N, 15.88. Found: C, 71.14; H, 7.18; N, 15.78. m.p.  $115-17^{\circ}\text{C}$

#### EXAMPLE 2

5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine



A solution of N-methyl, N-cyclohexyl 3-fluoro 4-bromomethyl benzamide (1.2 g 2.66 mm) and imidazopyridine (0.48 g 4.0 mm) in dimethylacetamide (25 ml) was heated overnight at 70-80°C with stirring under N<sub>2</sub>. Reaction solvent was removed in vacuo and the residue diluted with water and basified with ammonium hydroxide. The aqueous solution was extracted four times with chloroform and the combined organic extracts were backwashed three times with saturated aqueous sodium chloride solution. The organic solution was dried over magnesium sulfate, the drying agent was filtered and the filtrate concentrated in vacuo to give 0.88 g of the crude compound. Purification of the compound was effected by chromatography on silica using mixtures of chloroform, ethanol and ammonium hydroxide.


Analysis calcd for C <sub>21</sub> H <sub>23</sub> FN <sub>4</sub> O. 0.8 H <sub>2</sub> O:	C,66.22;	H,6.51;	N,14.71;	F,4.99.
Found:	C,66.03;	H,6.44;	N,14.65;	F,4.91.

mp 154-158°C

In the same manner as described in Example 2 the compounds of the Examples 3 to 11 described in Tables A & B were prepared.



TABLE A



Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	m.p. (°C)	Analysis	
								Calcd.	Found.
3			OCH <sub>3</sub>	1	H	H	217-19	C: 71.12	70.83
								H: 7.26	7.33
								N: 13.28	13.06
4			OCH <sub>3</sub>	1	H	H	197-99	C: 71.01	70.87
								H: 7.52	7.49
								N: 12.74	12.70
5			OCH <sub>3</sub>	1	H	H	192-95	C: 70.90	70.15
								H: 7.44	7.47
								N: 13.78	13.71
6		CH <sub>3</sub>	OCH <sub>3</sub>	1	H	H	206-8*	C: 69.15	68.78
								H: 6.86	6.88
								N: 14.61	14.57

TABLE A







Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	mpt. (°C)	Analysis Calcd. . . . Found.	Mol. form.
7			F	1	H	H	205-8	C: 71.4 H: 6.95 N: 13.33 F: 4.52	C <sub>25</sub> H <sub>29</sub> FN <sub>4</sub> O
8			F	1	H	H	178-82	C: 68.46 H: 6.99 N: 13.81 F: 4.71	C <sub>23</sub> H <sub>27</sub> FN <sub>4</sub> 0.5H <sub>2</sub> O
9		CH <sub>3</sub>	F	1	H	H	154-8	C: 66.22 H: 6.51 N: 14.71 F: 4.99	C <sub>21</sub> H <sub>23</sub> FN <sub>4</sub> 0.8H <sub>2</sub> O


TABLE B



Example	$R_1$	$R_2$	mpt. (°C)	Analysis		Mol. Formula
				Calcd	Found	
10		-CH <sub>3</sub>	167-67	C: 72.38 H: 6.14 N: 16.08	72.26 7.10 16.01	C <sub>21</sub> H <sub>24</sub> N <sub>4</sub> O
11			209-12	C: 73.51 H: 7.50 N: 14.88	72.97 7.63 14.61	C <sub>23</sub> H <sub>28</sub> N <sub>4</sub> O

EXAMPLE 12


5-[(4-(N,N-dicyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine



To a stirred solution of imidazopyridine (750 mg, 6.3 mmol) in N,N-dimethylacetamide, 4-bromomethyl-N,N-dicyclohexyl benzamide (2.6 g, 6.88 mmol) was added. The reaction mixture was stirred under argon at 80-85 °C. After 24h, the reaction flask was cooled to room temperature and the solvent removed under reduced pressure at < 45 °C. The residue obtained was triturated with ether (2 x 70 mL) and filtered. The crude (2.7 g) was chromatographed (silica gel, CH<sub>2</sub>Cl<sub>2</sub>-MeOH-NH<sub>4</sub>OH 80-20-1) to give pure product (1.47 g, 62%) which was recrystallized from EtOAc-CH<sub>3</sub>CN. mp 233-35 °C; Analysis calcd. for C<sub>26</sub>H<sub>32</sub>N<sub>4</sub>O. 0.3H<sub>2</sub>O: C, 74.0; H, 7.73; N, 13.28. Found C, 73.93; H, 7.90; N, 13.09.

In the same manner as described in Example 12 the compounds of the Examples 13 to 38 described in Table C were prepared.

TABLE C



Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	mp. (°C)	Analysis Calc.	Found	Mol. Formula
13		CH <sub>3</sub>	H	1	H	H	115-17	C: 71.46 H: 7.00 N: 15.88	71.14 7.18 15.79	C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> C <sub>21</sub> H <sub>24</sub> N <sub>2</sub>
14	1-(CH <sub>2</sub> ) <sub>7</sub> Cl	H	H	1	H	H	113-15	C: 72.52 H: 7.69 N: 15.38	72.26 7.72 15.28	C <sub>22</sub> H <sub>26</sub> N <sub>2</sub> C <sub>22</sub> H <sub>26</sub> N <sub>2</sub> C <sub>22</sub> H <sub>26</sub> N <sub>2</sub>
15	1-(CH <sub>2</sub> ) <sub>9</sub> Cl	H	H	1	H	H	141-45	C: 73.46 H: 8.16 N: 14.28	73.35 8.32 14.26	C <sub>24</sub> H <sub>28</sub> N <sub>2</sub> C <sub>24</sub> H <sub>28</sub> N <sub>2</sub> C <sub>24</sub> H <sub>28</sub> N <sub>2</sub>

TABLE C





Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	mp. (°C)	Analysis Calcd.	Found.	Mol. form.
16			H	1	H	H	209-10	C: 72.70 H: 7.48 N: 14.75	72.42 7.60 14.82	C <sub>23</sub> H <sub>28</sub> N <sub>2</sub>
17			H	1	H	H	210-11	C: 73.80 H: 7.69 N: 14.35	73.40 7.78 14.25	C <sub>24</sub> H <sub>30</sub> N <sub>2</sub>

TABLE C

Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	m.p. (°C)	Anal. Calcd.	Found	Mol. Formula
18	$-(CH_2)_{11}-$	H	H	1	H	H	150-2	C: 74.29 H: 8.57 N: 13.33	74.10 8.75 13.36	C <sub>26</sub> H <sub>36</sub> N <sub>2</sub> O
19		H	H	1	H	H	113-28	C: 73.31 H: 7.52 N: 13.68	73.41 7.79 13.33	C <sub>25</sub> H <sub>32</sub> N <sub>2</sub> O
20		H	H	1	H	H	221-2	C: 72.52 H: 7.69 N: 15.38	72.33 7.82 15.28	C <sub>22</sub> H <sub>28</sub> N <sub>2</sub> O

TABLE C







Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	mp <sub>lit</sub> (°C)	Analysis Calcd.	Found.	Mol. form.
21		CH <sub>3</sub>	H	2	H	H	223-5	C: 71.15 H: 7.27 N: 15.09	70.99 7.48 14.81	C <sub>22</sub> H <sub>26</sub> N <sub>2</sub> 0.5H <sub>2</sub> O
22		CH <sub>3</sub>	H	3	H	H		C: 70.86 H: 7.57 N: 14.36	70.12 7.64 14.23	C <sub>23</sub> H <sub>28</sub> N <sub>2</sub> 0.75H <sub>2</sub> O
23			H	1	H	CH <sub>3</sub>	147-98	C: 74.62 H: 7.46 N: 15.43	74.24 7.50 15.80	C <sub>25</sub> H <sub>30</sub> N <sub>2</sub>
24			H	1	CH <sub>3</sub>	H	225-8	C: 72.31 H: 7.48 N: 13.18	72.36 7.57 13.50	C <sub>25</sub> H <sub>30</sub> N <sub>2</sub> 0.7H <sub>2</sub> O



TABLE C









Example	R <sub>1</sub>	R <sub>2</sub>	X	R <sub>3</sub>	R <sub>4</sub>	m.p. (°C)	Analysis	
							Calcd.	Found.
25			H	H	H	190-3	C: 74.22 H: 7.21 N: 14.43	73.98 7.27 14.33
26			H	H	H	222-23	C: 74.62 H: 7.46 N: 13.93	74.21 7.45 14.26
27			H	H	H	233-35	C: 74.0 H: 7.73 N: 13.18	73.93 7.95 13.07
28			H	H	H	197-98	C: 73.84 H: 7.69 N: 14.35	73.86 7.87 14.35

TABLE C







Example.	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	m.p. (°C)	Analysis Calcd.	Found	Mol. Formula
29			H	1	H	H	213-14	C: 75.73 H: 6.16 N: 14.13	75.66 6.18 14.08	C <sub>15</sub> H <sub>17</sub> N
30			H	1	H	H	187-89	C: 74.94 H: 7.74 N: 13.44	74.58 7.84 13.32	C <sub>16</sub> H <sub>21</sub> N
31			H	1	H	H	211-12	C: 74.94 H: 7.74 N: 13.44	74.85 7.84 13.39	C <sub>17</sub> H <sub>23</sub> N

TABLE C















Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	m.p. (°C)	Anal. C, H, N	Found	Mol. Formula
32			H	1	H	H	198-200	C: 73.84 H: 7.69 N: 14.35	73.20 7.70 14.74	C <sub>24</sub> H <sub>30</sub> N <sub>2</sub>
33			H	1	H	H	211-13	C: 73.60 H: 7.35 N: 13.73	73.12 7.92 13.72	C <sub>25</sub> H <sub>32</sub> N <sub>2</sub>
34			H	1	H	H	88-90	C: 72.06 H: 7.04 N: 14.62	72.25 7.16 14.48	C <sub>23</sub> H <sub>25</sub> N <sub>2</sub>

TABLE C

Example	R <sub>1</sub>	R <sub>2</sub>	X	n	R <sub>3</sub>	R <sub>4</sub>	mpt. (°C)	Analysis Calcd. . . . .	Found.	Mol. form.
35			H	1	H	H	224-5	C: 71.42 H: 7.14 N: 16.66	71.30 7.29 16.71	C <sub>20</sub> H <sub>24</sub> N <sub>2</sub>
36			H	1	H	H	228-30	C: 74.22 H: 7.21 N: 14.43	74.04 7.29 14.28	C <sub>24</sub> H <sub>28</sub> N <sub>2</sub>
37		H	H	1	H	H	219-21	C: 70.33 H: 6.68 N: 16.41	70.34 6.87 16.28	C <sub>20</sub> H <sub>24</sub> N <sub>2</sub>
38			H	1	H	H	156-8	C: 71.85 H: 7.24 N: 15.24	71.85 7.24 15.14	C <sub>22</sub> H <sub>26</sub> N <sub>2</sub>

EXAMPLE 39

Preparation of 5-[4{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamide}benzyl]imidazo[4,5-c]pyridine




To a stirred solution of imidazopyridine (400 mg, 3.4 mmol) in N,N-dimethylacetamide (30 ml), 4-bromomethyl-N-cyclopentyl, N-3,5-dimethylcyclohexyl benzamide (1.4g, 3.57mmol) was added. The reaction mixture was stirred under argon at 80-85 °C. After 40h, the reaction flask was cooled to room temperature and the solvent removed under reduced pressure at <45 °C. The residue obtained was triturated with ether (2X70 ml) and filtered. The crude (1.8g) was chromatographed (silica gel, CH<sub>2</sub>Cl<sub>2</sub>-MeOH-NH<sub>4</sub>OH 90-10-1) to give pure product (1.05 g, 72%) which was recrystallized from EtOAc-CH<sub>3</sub>CN. mp 214-16 °C. Anal calcd. for C<sub>27</sub>H<sub>34</sub>N<sub>4</sub>O : C, 75.30; H, 7.9; N, 13.02. Found C, 74.92; H, 8.07; N, 12.97.

In the same manner as described in Example 39 the compounds of the Examples 40 to 55 described in Table D were prepared.

Table D

6116W



Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	M pt (°C)	Analysis Calcd.	Found	Molecular Formula
40			H	H	H	H	204-6	C 74.62 H 7.46 N 13.93	74.12 7.56 13.90	C <sub>25</sub> H <sub>30</sub> N <sub>4</sub> O
41			H	H	H	H	229-31	C 73.40 H 7.45 N 14.89	73.13 7.61 14.85	C <sub>23</sub> H <sub>28</sub> N <sub>4</sub> O
42			H	H	H	H	214-16	C 75.30 H 7.90 N 13.02	74.92 8.07 12.97	C <sub>27</sub> H <sub>34</sub> N <sub>4</sub> O
43			H	H	H	H	223-5	C 74.96 H 7.74 N 13.15	74.65 7.79 13.35	C <sub>26</sub> H <sub>32</sub> N <sub>4</sub> O
44			H	H	H	H	182-5	C 75.00 H 7.31 N 13.16	74.72 7.35 13.37	C <sub>26</sub> H <sub>30</sub> N <sub>4</sub> O 0.1H <sub>2</sub> O

Table D (cont'd.)










Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	M pt (°C)	Analysts Calcd.	Analysts Found	Molecular formula
45	H		H	H	H	H	242-4	C 66.79 H 5.26 N 13.97	66.79 4.97 13.26	C <sub>20</sub> H <sub>17</sub> N <sub>3</sub> O 0.9H <sub>2</sub> O
46			H	H	H	H	95-103	C 71.76 H 8.93 N 11.86	71.33 8.25 11.35	C <sub>25</sub> H <sub>32</sub> N <sub>4</sub> O 0.75H <sub>2</sub> O
47			OCH <sub>3</sub>	H	H	OH	235-37	C 68.22 H 7.13 N 13.26	67.66 7.23 13.09	C <sub>24</sub> H <sub>30</sub> N <sub>4</sub> O <sub>3</sub>
48			OCH <sub>3</sub>	H	H	Cl	171-3	C 65.22 H 6.62 N 12.70 Cl 8.10	64.96 6.78 12.31 8.47	C <sub>24</sub> H <sub>29</sub> N <sub>4</sub> O <sub>2</sub> Cl
49			OCH <sub>3</sub>	H	H	OCH <sub>3</sub>	212-14	C 66.71 H 7.50 N 12.90	66.53 7.25 12.26	C <sub>23</sub> H <sub>32</sub> N <sub>4</sub> O <sub>3</sub> 0.75H <sub>2</sub> O


Table D (cont'd.)

Example	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	M pt (°C)	Analysis Calcd. found	Molecular Formula
30			H	OCH <sub>3</sub>	H	H	236-8	C 72.60 72.28 H 7.67 7.65 N 12.54 12.46	C <sub>27</sub> H <sub>31</sub> N <sub>4</sub> O <sub>2</sub>
31			OCH <sub>3</sub>	H	H	H	186-8	C 72.60 72.21 H 7.67 7.91 N 12.54 12.28	C <sub>27</sub> H <sub>31</sub> N <sub>4</sub> O <sub>2</sub>
32			OCH <sub>3</sub>	OCH <sub>3</sub>	H	H	214-16	C 69.29 69.01 H 7.45 7.42 N 11.97 11.86	C <sub>27</sub> H <sub>31</sub> N <sub>4</sub> O <sub>3</sub> 0.3H <sub>2</sub> O
33			Br	OCH <sub>3</sub>	OCH <sub>3</sub>	H	191-3	C 57.95 57.56 H 6.30 6.01 N 10.01 9.93 Br 19.3 15.8	C <sub>27</sub> H <sub>31</sub> N <sub>4</sub> O <sub>3</sub> Br H <sub>2</sub> O
34			Br	H	OCH <sub>3</sub>	H	167-70	C 60.67 60.47 H 6.41 6.24 N 10.48 10.00 Br 14.95 14.57	C <sub>27</sub> H <sub>33</sub> N <sub>4</sub> O <sub>2</sub> Br 0.5H <sub>2</sub> O
35			OMe	H	H	H	210-13	C 64.41 64.40 H 7.09 7.14 N 12.51 12.41 Cl 7.92 8.0	C <sub>24</sub> H <sub>31</sub> N <sub>4</sub> ClO <sub>2</sub> 0.75H <sub>2</sub> O

EXAMPLE 56

Preparation of 5-[4{-(N-isopropyl,N-3-methylcyclopentyl)carboxamide}benzyl]imidazo[4,5-c]pyridine






To a stirred solution of imidazopyridine (689 mg, 5.76 mmol) in N,N-dimethylacetamide (30 ml), 4-bromomethyl-N-isopropyl,N-3-methylcyclohexyl benzamide (2.17g, 6.42mmol) was added. The reaction mixture was stirred under argon at 95 °C. After 48h, the reaction flask was cooled to room temperature and the solvent removed under reduced pressure at <45 °C. The residue obtained was triturated with ether (2X100ml) and filtered. The crude (2.97g) was chromatographed (silica gel, CH<sub>2</sub>Cl<sub>2</sub>-MeOH-NH<sub>4</sub>OH 90-10-1) to give pure product (.93g, 43%) which was recrystallized from EtOAc-CH<sub>3</sub>CN. mp 229-31 °C. Anal calcd. for C<sub>23</sub>H<sub>28</sub>N<sub>4</sub>O : C, 73.40; H, 7.45; N, 14.89. Found C, 73.13; H, 7.64; N, 14.85.

#### EXAMPLE 57


Preparation of 5-[4-((N-isopropyl,N-cyclohexyl)carboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine



To a stirred solution of imidazopyridine (1.5 g, 12.6 mmol) in dimethylacetamide (120 ml) under argon, N-isopropyl, N-cyclohexyl-3-methoxy-4-bromomethylbenzamide (5.1 g, 13.86 mmol) was added in one portion. The reaction temperature was slowly raised to 80-85 °C and was stirred over the week-end. The reaction flask was cooled to room temperature and the solvent removed under reduced pressure at <45 °C. The residue obtained was triturated with excess of dry ether (2X100 ml) and filtered. The crude product was chromatographed (silica gel; CH<sub>2</sub>Cl<sub>2</sub> : MeOH : NH<sub>4</sub>OH :: 90 : 10 : 1) to give pure alkylated product (3.53 g, 69%). The product could be recrystallized from ethyl acetate. mp 192-95 °C. Anal calcd. for C<sub>24</sub>H<sub>30</sub>N<sub>4</sub>O<sub>2</sub> : C, 70.90; H, 7.44; N, 13.78. Found C, 70.58; H, 7.43; N, 13.78.

#### EXAMPLE 58


Preparation of 5-[4-((N-isopropyl,N-cyclohexyl)carboxamido)-2-methoxybenzyl]-imidazo[4,5-c]pyridine hydrochloride



To clear solution of the product of Example 57 (100 mg) in methanol (7 ml), HCl in dioxane (5ml, 6N solution) was added. After stirring at room temp. for 2h, the solvent was removed under reduced pressure. Ethyl acetate (25ml) was added and mixture was refluxed for 1h. The contents were filtered hot and the residue was washed with more hot ethyl acetate. After drying, the product (92mg) was collected, mp 210-13 °C. Anal calcd. for  $C_{24}H_{31}N_4ClO_2 \cdot 0.25H_2O$  : C, 64.41; H, 7.09; N, 12.52; Cl, 7.92. Found C, 64.40; H, 7.34; N, 12.43, Cl, 8.0.

#### EXAMPLE 59


Preparation of 5-[4-(N-cyclopentyl, N-3-methylcyclohexyl)carboxamide]-3-methoxybenzyl]imidazo[4,5-c]pyridine



To a stirred solution of imidazopyridine (412mg, 3.47mmol) in N,N-dimethylacetamide (25 ml), 4-bromomethyl-2-methoxy-N-cyclopentyl, N-3-methylcyclohexyl benzamide (1.49g, 3.65 mmol) was added. The reaction mixture was stirred under argon at 90-95 °C. After 48h, the reaction flask was cooled to room temperature and the solvent removed under reduced pressure at <45 °C. The residue obtained was triturated with ether (2X70ml) and filtered. The crude (1.85g) was chromatographed (silica gel,  $CH_2Cl_2$ -MeOH-NH<sub>4</sub>OH 90-10-1) to give pure product (1.05g, 67%) which was recrystallized from EtOAc. mp 226-28 °C. Anal calcd. for  $C_{27}H_{34}N_4O_2$  : C, 72.60; H, 7.67; N, 12.54. Found C, 72.28; H, 7.65; N, 12.44.

#### EXAMPLE 60


Preparation of 5-[4-(N-cyclopentyl, N-3-methylcyclohexyl)carboxamido]-2-methoxybenzyl]imidazo[4,5-c]pyridine



To a stirred solution of imidazopyridine (525 mg, 4.4 mmol) in N,N-dimethylacetamide (25 ml), 4-bromomethyl-3-methoxy-N-cyclopentyl, N-3-methylcyclohexyl benzamide (1.9g, 4.66mmol) was added. The reaction mixture was stirred under argon at 90-95 °C. After 48h, the reaction flask was cooled to room temperature and the solvent removed under reduce dpressure at <45 °C. The residue obtained was triturated with ether (2X100ml) and filtered. The crude was chromatographed (silica gel, CH<sub>2</sub>Cl<sub>2</sub>-MeOH-NH<sub>4</sub>OH 90-10-1) to give pure product (1.39 g, 71%) which was recrystallized from EtOAc-CH<sub>3</sub>CN, mp 186-88 °C. Anal calcd. for C<sub>27</sub>H<sub>34</sub>N<sub>4</sub>O<sub>2</sub> : C, 72.60; H, 7.67; N, 12.54. Found C, 72.21; H, 7.91; N, 12.28.

#### EXAMPLE 61


Preparation of 5-[4-(N-cyclopentyl,N-cyclohexyl)carboxamido]-2-methoxybenzyl]imidazo[4,5-c]pyridine



To a stirred solution of imidazopyridine (660mg, 5.57 mmol) in N,N-dimethylacetamide (25ml), 4-bromomethyl-2-methoxy-N-cyclopentyl,N-cyclohexyl benzamide (2.0g, 5.07mmol) was added. The reaction mixture was stirred under argon at 75 °C. After 24h, the reaction flask was cooled to room temperature and the solvent removed under reduced pressure. The residue obtained was diluted with water (650ml) and basified with aq. ammonium hydroxide (20ml). The reaction solution was extracted with chloroform (4x100 ml). The organic layer was washed with brine (3x250 ml), dried (MgSO<sub>4</sub>) and filtered. The combined filtrate was concentrated and the residue (2.79g) chromatographed (silica gel, CHCl<sub>3</sub>-EtOH-NH<sub>4</sub>OH 10-90-1) to give desired product (1.36g, 62%), mp 197-99 °C. Anal calcd. for C<sub>26</sub>H<sub>32</sub>N<sub>4</sub>O<sub>2</sub>·0.4H<sub>2</sub>O : C, 71.01; H, 7.52; N, 12.74. Found C, 70.87; H, 7.49; N, 12.70.

#### EXAMPLE 62

Preparation of 5-[4-(N-isopropyl,N-cyclohexyl) carboxamide]benzyl]imidazo[4,5-c]pyridine



5

To a stirred solution of imidazopyridine (680mg, 5.8mmol) in N,N-dimethylacetamide (30ml), 4-bromomethyl-N-isopropyl,N-cyclohexyl benzamide (2.2g, 6.44mmol) was added. The reaction mixture was stirred under argon at 80-85°C. After 20h, the reaction flask was cooled to room temperature and the solvent removed under reduced pressure at <45°C. The residue obtained was triturated with ether and filtered. The crude (1.85g) was chromatographed (silica gel, CH<sub>2</sub>Cl<sub>2</sub>-EtOH-NH<sub>4</sub>OH 80-20-1) to give pure product (1.29 g, 59%) which was recrystallized from EtOAc-CH<sub>3</sub>CN, mp 209-10°C. Anal calcd. for C<sub>23</sub>H<sub>28</sub>N<sub>4</sub>O 0.2H<sub>2</sub>O: C, 72.70; H, 7.48; N, 14.75. Found C, 72.92; H, 7.60; N, 14.82.

15


EXAMPLE 63

20

Preparation of 5-[4-[(N-cyclohexyl,N-isopropyl)carboxamide]-2-methoxybenzyl]-4-chloro-imidazo[4,5-c]pyridine

25

30



35

Preparation of the 4-chloro-imidazo [4,5-c]pyridine starting material as well as the 2-methoxy-3-bromobenz(N-cyclohexyl,N-isopropyl)amide have been described earlier in this specification. Coupling of the 4-chloro-imidazo[4,5-c]to the 2-methoxy-3-bromobenz-(N-cyclohexyl,N-isopropyl)amide in dimethylacetamide at 85-90°C for 26h gives the titled compound.


45

EXAMPLE 64


50

Preparation of


55



10 and



25 The above compounds can be synthesized according to the following scheme -



The N-1 compound of imidazopyridine is protected by SEM-C1 and converted to a pyridine N-oxide using m-chloroperbenzoic acid in a manner described for the preparation of 4-chloro-imidazo[4,5-c]pyridine. The pyridine-N-oxide compound is refluxed in acetic anhydride for 4 hrs. to give 4-oxo-1-(2-trimethylsilyl)-ethoxymethyl-imidazo[4,5-c]pyridine. Reacting this compound with 4-bromomethyl-3-methoxy-benz-[N-isopropyl,N-cyclohexyl) amide in dimethylformamide and sodium hydride at room temperature for 4 hours gives the 5-benzylated product. Cleavage of the SEM-group is accomplished by trifluoroacetic acid at 50 °C for 18 hours to give the compound of formula 8 (titled compound). Treatment of the 4-hydroxy group of the compound of formula 8 with sodium hydride/iodomethane gives the compound of formula 9 (titled compound).

#### EXAMPLE 65

##### PAF-induced platelet aggregation and secretion:

Washed, [<sup>3</sup>H]serotonin-labeled rabbit platelets were prepared as previously described in COX, C. P., J. LINDEN and S. I. SAID: VIP elevates platelet cyclic AMP (cAMP) levels and inhibits *in vitro* platelet activation induced by platelet-activating factor (PAF). Peptides 5: 25-28, 1984, and maintained in an atmosphere of 5% CO<sub>2</sub> at 37 °C until used in the bioassay. Aliquots of platelets (2.5 x 10<sup>8</sup>/ml) were incubated with either an antagonist of PAF or the appropriate vehicle for 60 sec prior to the addition of PAF (0.1 nM to 0.2 μM). Aggregation was continuously monitored on a strip-chart recorder and recorded as the height of the tracing at 60 sec after the addition of PAF. Secretion of [<sup>3</sup>H] serotonin was measured in a sample of the platelet suspension removed at 60 sec after the addition of PAF. The percent inhibition of aggregation and secretion was calculated by comparing antagonist-treated platelets with the appropriate vehicle-treated control platelets. Each combination of antagonist and PAF was repeated 12-15 times, using several different platelet preparations. IC<sub>50</sub> values were determined by inspection of the dose-response curves.

#### EXAMPLE 66

##### Inhibition of <sup>3</sup>H-PAF Binding to Human Platelet Membrane Receptors

##### Receptor Preparation:

Ten units of in-dated human packed platelets, each containing 45-65 ml platelet rich-plasma, were purchased from a commercial blood bank. Disposable plasticware was used throughout for receptor preparation. The units were pooled and a 1 ml aliquot was removed for determination of platelet concentration, using a Coulter Counter. The remaining platelet rich plasma was dispensed into 50 ml conical tubes and centrifuged at room temperature for 15 minutes at 3000 RPM (2300 x g). Plasma was decanted and the platelets were resuspended in 35 ml of buffer (10 mM Trizma 7.0, 2 mM EDTA (dipotassium salt), and 150 mM KCl) and transferred to fresh tubes, which were centrifuged again as above. The platelets were washed 3 times, avoiding contaminating erythrocytes at the bottom of the pellets. Pellets were consolidated at each step, and by the last wash with EDTA/KCl buffer, most of the erythrocytes were in 1 tube. The pellets were resuspended in buffer containing 10 mM Trizma 7.0 with 10 mM CaCl<sub>2</sub>. Following centrifugation, the buffer was decanted and the pellets were resuspended in the CaCl<sub>2</sub> buffer, avoiding erythrocyte contamination by recovering less than 100% of the platelet pellets. The resuspended platelets were dispensed in 8-10 ml aliquots into Corex tubes and disrupted by three cycles of freezing (dry ice/ethanol) and thawing (24 °C). The tubes were centrifuged at 40,000 x g for 20 minutes at 4 °C. Supernatants were decanted and each pellet was resuspended in 5-7 ml 10 mM Trizma 7.0. All resuspended pellets were pooled and aliquots of about 1200 μl were dispensed into 1.5 ml microfuge tubes and frozen at -70 °C. Protein content was determined by a fluorescamine protein assay.

Assay Methods: Receptor Characterization

Each receptor preparation was evaluated to determine the number of receptor populations, the number of PAF receptor equivalents/mg protein and the dissociation constant ( $K_D$ ) for PAF binding. This required 2-3 experiments in which the protein concentration was held constant and the  $^3\text{H}$ -PAF ligand concentration was varied from approximately 0.10-2.5 nM and the data was analyzed by Scatchard methodology. Total incubation volume was 250  $\mu\text{l}$  for these procedures and incubations were conducted at 24 °C for 30 minutes. For further experimentation, total incubation volumes are 500  $\mu\text{l}$ . Protein and ligand concentrations were adjusted to give 0.075 nM receptor equivalents in the presence of 0.75 nM  $^3\text{H}$ -PAF. Each receptor preparation was then used to determine the dose - response displacement relationship of unlabeled PAF and the PAF antagonist, triazolam. As long as the  $K_D$  value and  $\text{IC}_{50}$  values for PAF and triazolam were consistent with similar data collected from past receptor preparations used in the assay, the new receptor preparation was used for evaluating compounds.

Assay Methods: Routine Assay of Compounds

The compounds were weighed precisely and solubilized in quantities of DMSO such that a 5  $\mu\text{l}$  aliquot in the incubate would deliver the desired compound concentration. Compounds tested for the first time in this assay were evaluated at a concentration of 50  $\mu\text{M}$  in the incubation medium. All compounds were generally solubilized in DMSO for about 2 hours prior to assay. Triazolam was always included in each screening assay as a compound inhibition control. A standard concentration of 50  $\mu\text{M}$  inhibited  $^3\text{H}$ -PAF binding by approximately 50%. Nonspecific binding control solution was made by drying to completion about 26.2  $\mu\text{l}$  unlabeled PAF under a stream of argon. PAF was resolubilized in 1000  $\mu\text{l}$  DMSO. When delivered in a 5  $\mu\text{l}$  aliquot, the final concentration of 1  $\mu\text{M}$  PAF in the incubate exceeded by 1000-fold the concentration of  $^3\text{H}$ -PAF.

All buffers containing proteins were made at room temperature on the day of assay. Assay buffer was prepared by adding 125 mg human albumin to 25 ml of stock buffer (10 mM Trizma 7.4 with 20 mM  $\text{CaCl}_2$ ). Rinse buffer was made by adding 20 grams bovine serum albumin to 1000 ml stock buffer. About 80 ml of rinse buffer was decanted into a small pyrex dish and used to soak 65 Whatman GF/C 2.5 cm glass filters. The remaining rinse buffer was poured into a repipet and placed into an ice bath along with the filters.

Ligand for assay was prepared by adding about 10  $\mu\text{l}$  of stock  $^3\text{H}$ -PAF (DuPont NEN, NET-668) to 14 ml of assay buffer. Since the amount of  $^3\text{H}$ -PAF in the final incubate was to be 0.75 nM, the actual amount of stock  $^3\text{H}$ -PAF to be used had to be determined for each lot of material based upon its specific activity.

Membrane receptors for assay were prepared by thawing the appropriate number of tubes at room temperature and adding membranes to 10 mM Trizma 7.0 containing 10 mM  $\text{CaCl}_2$ . A total volume of 14 ml was made. The actual amount of membranes needed was determined by the requirement to have 0.075 nM PAF receptor equivalents per assay tube. All materials were kept in motion by rocking on a rocker plate.

First, 5  $\mu\text{l}$  of compound of DMSO was added to each 12 x 75 mm polypropylene tube, followed by the addition of 95  $\mu\text{l}$  assay buffer. Next, 200  $\mu\text{l}$   $^3\text{H}$ -PAF was added to each tube and 3 aliquots of  $^3\text{H}$ -PAF taken at different times during the dispensing were placed in scintillation vials. The reaction was initiated by the addition of 200  $\mu\text{l}$  of membranes. All tubes were very briefly vortexed and placed in a 24 °C water bath for about 30 minutes. During this time, Whatman GF/C filters were placed on the filter racks of 5 Millipore vacuum manifolds. The incubations were terminated by first adding 4 ml ice-cold rinse buffer to each incubation tube and then decanting them over the filters under vacuum. Tubes and filters were rinsed twice more. Each filter was placed into a 20 ml scintillation vial to which 20 ml Aquasol (DuPont NEN, NDF 952) was added. All vials were given 2 hours in the dark for photo and chemiluminescence to dissipate prior to liquid scintillation counting.

In summary, each incubation tube contained 500  $\mu\text{l}$  total volume of incubate. This consisted of 5  $\mu\text{l}$  drug with DMSO or only DMSO, 95  $\mu\text{l}$  assay buffer, 200  $\mu\text{l}$   $^3\text{H}$ -PAF (0.75 nM final concentration) and 200 microliters membrane receptors (0.075 nM final concentration). 60 tubes per assay were run and each dose was performed in triplicate. Controls in every assay consisted of 2 diluent (DMSO) "0" controls (2 triplicate determinations placed at different positions within the 60 tube assay), 1 nonspecific binding control, and 1 triazolam drug control. The 16 remaining doses were used to test 16 different compounds at the screening dose of 50  $\mu\text{M}$ , or to run dose-response determinations for a compound. In general, dose-response curves were composed of 4 compound doses designed to inhibit  $^3\text{H}$ -PAF binding by 15-85%, with at least 1 dose on each side of the 50% point.

Routine Assay Calculations:

5 Triplicate DPM determinations (corrected for background) within a single compound dose were averaged while all 6 determinations of total binding ("0" dose, DMSO only) were averaged. The amount for nonspecific binding (1  $\mu$ M PAF) was subtracted from all the dose averages, giving an amount of specific binding in all cases. The percent displacement of  $^3$ H-PAF or inhibition of binding was calculated by the formula  $STBo - SBC / STBo \times 100$ , where STBo = specific binding of "0" dose controls and SBC = specific binding in the presence of compound. If a compound tested at the initial screening dose of 50  $\mu$ M inhibited binding by 45% or more, the compound was considered active and was tested in a dose-response manner  
10 to determine an  $IC_{50}$  value.

Compounds inhibiting PAF binding by less than 45% at a 50  $\mu$ M concentration were considered inactive and no further testing was done.

15  $IC_{50}$  values were determined on active compounds in subsequent tests. Three or more compound doses must inhibit  $^3$ H-PAF binding between 15-85%. Using a computer program, % displacement data was transformed (logit) and a least squares linear regression was performed on the data meeting the 15-85% requirement to determine  $IC_{50}$  values from data points derived from the same assay.



	Compound	PAF induced platelet secretion (IC <sub>50</sub> )M	PAF induced platelet aggregation (IC <sub>50</sub> )M	Inhibition of <sup>3</sup> H-PAF Binding to Human Platelet (IC <sub>50</sub> )μM
5				
	5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	7.2 x 10 <sup>-7</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	15.2
10				
	5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	11.0
	5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	9.71
15				
	5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]pyridine	1 to 5 x 10 <sup>-7</sup>	10 <sup>-6</sup> to 10 <sup>-7</sup>	11.9
	5-[4-(N-2-decyl-1-N-methylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-6</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	13.2
20				
	5-[4-(N-2-(4,4-dimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-6</sup>	10 <sup>-5</sup>	22.3
25				
	5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-7</sup> to 10 <sup>-8</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	7.65
	5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 5 x 10 <sup>-8</sup>	0.31
30				
	5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-5</sup>	19.3
	5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-7</sup> to 10 <sup>-6</sup>	10 <sup>-6</sup> to 10 <sup>-5</sup>	5.20
35				
	5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.17
40				
	5-[4-(N-sec-butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 5 x 10 <sup>-8</sup>	0.58
	5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-7</sup>	10 <sup>-6</sup>	2.82
45				
50				
55				

	Compound	PAF induced platelet secretion (IC <sub>50</sub> )M	PAF induced platelet aggregation (IC <sub>50</sub> )M	Inhibition of <sup>3</sup> H-PAF Binding to Human Platelet (IC <sub>50</sub> )μM
5	5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-7</sup> to 10 <sup>-8</sup>	10 <sup>-6</sup> to 10 <sup>-7</sup>	—
10	5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>		3.68
15	5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.0199
20	5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.32
25	5-[4-(N,N-dicyclohexylcarboxamidobenzyl)imidazo[4,5-c]pyridine	10 <sup>-8</sup>	10 <sup>-6</sup> to 10 <sup>-7</sup>	1.06
30	5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine	10 <sup>-5</sup> to 10 <sup>-6</sup>	10 <sup>-4</sup> to 10 <sup>-5</sup>	—
	5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine	10 <sup>-5</sup> to 10 <sup>-6</sup>	10 <sup>-4</sup> to 10 <sup>-5</sup>	61.1
35	5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.055
40	5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-8</sup>	0.0302
45	5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine	10 <sup>-8</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.0665


50

55

	Compound	PAF induced platelet secretion (IC <sub>50</sub> )M	PAF induced platelet aggregation (IC <sub>50</sub> )M	Inhibition of <sup>3</sup> H-PAF binding to Human Platelet (IC <sub>50</sub> )μM
5				
10	5-[4-(N-methyl-N-cyclohexyl carboxamido)-2-methoxybenzyl] imidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	—
	5-[4-(N-cyclopentyl-N- cyclohexylcarboxamido)-2- fluorobenzyl]imidazo [4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.0755
15	5-[4-(N-isopropyl-N-cyclohexyl carboxamido)-2-fluorobenzyl] imidazo[4,5-c]pyridine	10 <sup>-7</sup> to 10 <sup>-8</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.442
20	5-[4-(N-methyl-N-cyclohexyl carboxamido)-2-fluorobenzyl] imidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	—
	5-[4-(N-tert.butyl-N- cyclohexylcarboxamido) benzyl]imidazo[4,5- c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	87.7% inhib (50μM)
25	5-[4-(N-phenyl-N- cyclopentylcarboxamido) benzyl]imidazo[4,5- c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-6</sup>	2.35
30	5-[4-(N-1-methylcyclohexyl-N- cyclopentylcarboxamido)benzyl] imidazo[4,5-c]pyridine	10 <sup>-8</sup> to 10 <sup>-9</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	~ 0.074
	5-[4-(N-4-methylcyclohexyl-N- cyclopentylcarboxamido)benzyl] imidazo[4,5-c]pyridine	10 <sup>-7</sup> to 10 <sup>-8</sup>	10 <sup>-6</sup> to 10 <sup>-7</sup>	0.75
35	5-[3-(N-methyl-N-cyclohexyl carboxamido)benzyl]imidazo[4,5- c]pyridine	10 <sup>-4</sup> to 10 <sup>-5</sup>	10 <sup>-4</sup> to 10 <sup>-5</sup>	38% inhib (50μM)
	5-[3-(N-isopropyl-N-cyclohexyl carboxamido)benzyl]imidazo[4,5- c]pyridine	10 <sup>-5</sup> to 10 <sup>-6</sup>	10 <sup>-4</sup> to 5 x 10 <sup>-5</sup>	26.1% inhib (50μM)
40	5-[4-(N,N-dicyclopentyl carboxamido)benzyl]- 4-methylimidazo[4,5-c]pyridine	10 <sup>-7</sup> to 10 <sup>-8</sup>	10 <sup>-7</sup> to 10 <sup>-8</sup>	0.188
45	5-[4-(N,N-dicyclopentyl carboxamido)benzyl]- 2-methylimidazo[4,5-c]pyridine	10 <sup>-6</sup> to 10 <sup>-7</sup>	10 <sup>-5</sup> to 10 <sup>-6</sup>	70.8% inhib (50μM)
50				
55				

## Claims

1. A compound of the formula



or a pharmaceutically acceptable acid addition thereof: wherein

$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; heterocyclical-  
 20 heteroaromatic having 5 or 6 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen


$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or  
 30 branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxyalkyl wherein the alkyl groups are each 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl group are each 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms; and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.

$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

2. A compound according to Claim 1 having the formula



or a pharmaceutically acceptable acid addition salt thereof: wherein

$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso

that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen;

$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 position of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

3. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)-carboxamido}benzyl]imidazo[4,5-c]pyridine.

4. A compound according to Claim 2 which is 5-[4{-(N-isopropyl,N-3-methylcyclopentyl)-carboxamido}benzyl]imidazo[4,5-c]pyridine.

5. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)-carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine.

6. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)-carboxyamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.

7. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl-N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.

8. A compound according to Claim 2 which is 5-[4{-(N-isopropyl,N-cyclohexyl) carboxamido}benzyl]imidazo[4,5-c]pyridine.

9. A compound according to Claim 2 wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl.

10. A compound according to Claim 9 wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; or cycloalkyl having 3 to 8 carbon atoms.

11. A compound according to Claim 2 where  $Y$  is phenyl.


12. A compound according to Claim 2 wherein  $Y$  is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms.

13. A compound according to Claim 12 wherein the halogen is fluoro.

14. A compound according to Claim 12 wherein the alkoxy is methoxy.

15. A compound according to Claim 2 wherein  $n$  is an integer of 1 to 3.

16. A compound according to Claim 1 having the formula



or a pharmaceutically acceptable acid addition salt thereof: wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl.  $n$  is an integer of 1 to 3.  $R_3$  is hydrogen or alkyl of 1 to 6 carbon atoms and  $R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

17. A compound according to Claim 16 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

18. A compound according to Claim 16 which is 5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

19. A compound according to Claim 16 which is 5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]-pyridine.
20. A compound according to Claim 16 which is 5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]-pyridine.
21. A compound according to Claim 16 which is 5-[4-(N-2-decalyl)-N-methylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
22. A compound according to Claim 16 which is 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
23. A compound according to Claim 16 which is 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
24. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
25. A compound according to Claim 16 which is 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
26. A compound according to Claim 16 which is 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
27. A compound according to Claim 16 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
28. A compound according to Claim 16 which is 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
29. A compound according to Claim 16 which is 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
30. A compound according to Claim 16 which is 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
31. A compound according to Claim 16 which is 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
32. A compound according to Claim 16 which is 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
33. A compound according to Claim 16 which is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
34. A compound according to Claim 16 which is 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)]imidazo[4,5-c]pyridine.
35. A compound according to Claim 16 which is 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]-ethyl]imidazo[4,5-c]pyridine.
36. A compound according to Claim 16 which is 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]-propyl]imidazo[4,5-c]pyridine.
37. A compound according to Claim 16 which is 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
38. A compound according to Claim 16 which is 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
39. A compound according to Claim 16 which is 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
40. A compound according to Claim 16 which is 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
41. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine.
42. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine.
43. A compound according to Claim 16 which is 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
44. A compound according to Claim 16 which is 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
45. A compound according to Claim 1 wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring or phenyl. Y is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms. n is an integer of 1 to 3. R<sub>3</sub> is hydrogen or alkyl of 1 to 6 carbon atoms and R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms.

46. A compound according to Claim 45 wherein the halogen is fluoro.
47. A compound according to Claim 45 which is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
48. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
49. A compound according to Claim 45 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
50. A compound according to Claim 45 wherein the alkoxy is methoxy.
51. A compound according to Claim 45 which is 5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
52. A compound according to Claim 45 which is 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
53. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
54. A compound according to Claim 45 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
55. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride.
56. A pharmaceutical composition useful for treating diseases or disorder mediated by platelet-activating factor comprising at least one compound according to Claim 1, together with one or more non-toxic pharmaceutically acceptable carriers.
57. A pharmaceutical composition according to Claim 56 wherein said compound is selected from the group consisting of
- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-n-octylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-n-decylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-n-dodecylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-2-decyl-N-methylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine
  - 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclopentylcarboxamido)-2-methylbenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride
  - 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
  - 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine

5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido]benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl,N-3-methylcyclopentyl)carboxamido]benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido]-3-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido]-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl,N-cyclohexyl)carboxamido]-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl,N-cyclohexyl)carboxamido]benzyl]imidazo[4,5-c]pyridine

58. A method for treating diseases or disorder mediated by platelet-activating factor comprising administering a therapeutically effective dose of at least one compound of Claim 1 to a mammal in need of such treatment.

59. A method according to Claim 58 wherein said compound is selected from the group consisting of


5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-2-decyl-N-methylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-sec-butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclohexylcarboxamido)imidazo[4,5-c]pyridine  
 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine  
 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine  
 5-[4-(N-tert-butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-phenyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-3-N-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-4--methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride  
 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine.  
 5-[4-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido]benzyl]imidazo[4,5-c]pyridine.  
 5-[4-(N-isopropyl,N-3-methylcyclopentyl)carboxamido]benzyl]imidazo[4,5-c]pyridine.  
 5-[4-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido]-3-methoxybenzyl]imidazo[4,5-c]pyridine.  
 5-[4-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido]-2-methoxybenzyl]imidazo[4,5-c]pyridine.  
 5-[4-(N-cyclopentyl,N-cyclohexyl)carboxamido]-2-methoxybenzyl]imidazo[4,5-c]pyridine.  
 5-[4-(N-isopropyl,N-cyclohexyl)carboxamido]benzyl]imidazo[4,5-c]pyridine.



60. A method for treating diseases or disorder mediated by platelet-activating factor comprising administering a therapeutically effective dose of pharmaceutical composition of Claim 56 to mammal in need of such treatment.

Claims for the following Contracting State: ES

1. Process for preparing a compound of the formula



or a pharmaceutically acceptable acid addition thereof: wherein


$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; heterocyclical-  
 25 heteroaromatic having 5 or 6 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxyalkyl wherein the alkyl groups are each 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms, alkylthioalkyl wherein the alkyl group are each 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms; and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.


$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms,

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms, characterized in that an imidazopyridine of the formula II




wherein  $R_3$  and  $R_4$  are defined as hereinabove is reacted with a haloalkylbenzamide of the general formula (III)



wherein  $R_1$  and  $R_2$  and  $n$  are defined as before and  $X$  represents chloro, bromo, or methanesulfonyloxy and wherein the phenyl moiety may be substituted once or more by halogen, alkyl of 1 to 6 carbon atoms; alkoxy wherein the alkyl is 1 to 6 carbon atoms; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxy alkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl is 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms, thus yielding the desired final compounds of formula (I) which can optionally be transformed into the desired pharmaceutically acceptable acid addition salts by contacting said compounds of formula (I) with an appropriate acid.

2. A process according to Claim 1 wherein the compound prepared has the formula



or a pharmaceutically acceptable acid addition salt thereof; wherein

$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen;

$n$  is an integer of 1 to 5,

$R_3$  is a group substituted at one or more of the 4, 6, or 7 position of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms,

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.


3. A process according to Claim 2 wherein the compound prepared is

5-[4{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine.

4. A process according to Claim 2 wherein the compound prepared is

5-[4{-(N-isopropyl,N-3-methylcyclopentyl)carboxamido}benzyl]imidazo[4,5-c]pyridine.

5. A process according to Claim 2 wherein the compound prepared is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine.
6. A process according to Claim 2 wherein the compound prepared is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxyamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.
7. A process according to Claim 2 wherein the compound prepared is 5-[4{-(N-cyclopentyl,N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.
8. A process according to Claim 2 wherein the compound prepared is 5-[4{-(N-isopropyl,N-cyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine.
9. A process according to Claim 2 wherein  $R_1$  and  $R_2$  of the compound prepared are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl.
10. A process according to Claim 9 wherein  $R_1$  and  $R_2$  of the compound prepared are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; or cycloalkyl having 3 to 8 carbon atoms.
11. A process according to Claim 2 where Y is phenyl.
12. A process according to Claim 2 wherein Y is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms.
13. A process according to Claim 12 wherein the halogen is fluoro.
14. A process according to Claim 12 wherein the alkoxy is methoxy.
15. A process according to Claim 2 wherein n is an integer of 1 to 3.
16. A process according to Claim 1 wherein the compound prepared having the formula



- or a pharmaceutically acceptable acid addition salt thereof: wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl. n is an integer of 1 to 3.  $R_3$  is hydrogen or alkyl of 1 to 6 carbon atoms and  $R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.
17. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
18. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
19. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
20. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
21. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-2-decalyl)-N-methylcarboxamido]benzyl]imidazo[4,5-c]pyridine.
22. A process according to Claim 16 wherein the compounds prepared is 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
23. A process according to Claim 16 wherein the compound prepared is 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
24. A process according to Claim 16 wherein the compound prepared is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
25. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
26. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

27. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
28. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
29. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
30. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
31. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
32. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
33. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
34. A process according to Claim 16 wherein the compound prepared is 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)imidazo[4,5-c]pyridine.
35. A process according to Claim 16 wherein the compound prepared is 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine.
36. A process according to Claim 16 wherein the compound prepared is 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine.
37. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
38. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
39. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
40. A process according to Claim 16 wherein the compound prepared is 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
41. A process according to Claim 16 wherein the compound prepared is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine.
42. A process according to Claim 16 wherein the compound prepared is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine.
43. A process according to Claim 16 wherein the compound prepared is 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
44. A process according to Claim 16 wherein the compound prepared is 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
45. A process according to Claim 1 wherein  $R_1$  and  $R_2$  of the compound prepared are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring or phenyl. Y is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms. n is an integer of 1 to 3.  $R_3$  is hydrogen or alkyl of 1 to 6 carbon atoms and  $R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.
46. A process according to Claim 45 wherein the halogen is fluoro.
47. A process according to Claim 45 wherein the compound prepared is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
48. A process according to Claim 45 wherein the compound prepared is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
49. A process according to Claim 45 wherein the compound prepared is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
50. A process according to Claim 45 wherein the compound prepared is is methoxy.
51. A process according to Claim 45 wherein the compound prepared is 5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
52. A process according to Claim 45 wherein the compound prepared is 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
53. A process according to Claim 45 wherein the compound prepared is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
54. A process according to Claim 45 wherein the compound prepared is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.

55. A process according to Claim 45 wherein the compound prepared is  
5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride.

56. Use of at least one compound according to Claim 1 for preparing a medicament for treating diseases or disorder mediated by platelet-activating factor.

57. Use according to Claim 56 wherein said compound is selected from the group consisting of


- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-n-octylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-n-decylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-n-dodecylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-2-decyl-N-methylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)imidazo[4,5-c]pyridine
- 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine
- 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine
- 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)-2-methylbenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride
- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
- 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine
- 5-[4-{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-isopropyl,N-3-methylcyclopentyl)carboxamido}benzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-isopropyl,N-cyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine

50

Claims for the following Contracting State: GR

1. A compound of the formula

55



or a pharmaceutically acceptable acid addition thereof: wherein

$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; heterocyclical-  
 15 alkyl having 4 to 8 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; heteroaromatic having 5 or 6 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with  
 20 the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen


$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or  
 25 branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxyalkyl wherein the alkyl groups are each 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl group are each 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl  
 30 is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms; and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.

$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is  
 35 selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

2. A compound according to Claim 1 having the formula



or a pharmaceutically acceptable acid addition salt thereof: wherein

$R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; phenyl;  
 50 substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso  
 55 that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the

phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen;

5 n is an integer of 1 to 5.

R<sub>3</sub> is a group substituted at one or more of the 4, 6, or 7 position of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms.

10 3. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)-carboxamido}benzyl]imidazo[4,5-c]pyridine.

4. A compound according to Claim 2 which is 5-[4{-(N-isopropyl,N-3-methylcyclopentyl)-carboxamido}benzyl]imidazo[4,5-c]pyridine.

15 5. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)-carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine.

6. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)-carboxyamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.

7. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl-N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.

20 8. A compound according to Claim 2 which is 5-[4{-(N-isopropyl,N-cyclohexyl) carboxamido}benzyl]-imidazo[4,5-c]pyridine.

9. A compound according to Claim 2 wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl.

25 10. A compound according to Claim 9 wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; or cycloalkyl having 3 to 8 carbon atoms.

11. A compound according to Claim 2 where Y is phenyl.


30 12. A compound according to Claim 2 wherein Y is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms.

13. A compound according to Claim 12 wherein the halogen is fluoro.

14. A compound according to Claim 12 wherein the alkoxy is methoxy.

15. A compound according to Claim 2 wherein n is an integer of 1 to 3.

35 16. A compound according to Claim 1 having the formula



or a pharmaceutically acceptable acid addition salt thereof: wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl. n is an integer of 1 to 3. R<sub>3</sub> is hydrogen or alkyl of 1 to 6 carbon atoms and R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms.

50 17. A compound according to Claim 16 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.

18. A compound according to Claim 16 which is 5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

55 19. A compound according to Claim 16 which is 5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

20. A compound according to Claim 16 which is 5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

21. A compound according to Claim 16 which is 5-[4-(N-2-decalyl)-N-methylcarboxamido]benzyl]-imidazo[4,5-c]pyridine.
22. A compound according to Claim 16 which is 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
23. A compound according to Claim 16 which is 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
24. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
25. A compound according to Claim 16 which is 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
26. A compound according to Claim 16 which is 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
27. A compound according to Claim 16 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
28. A compound according to Claim 16 which is 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
29. A compound according to Claim 16 which is 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
30. A compound according to Claim 16 which is 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
31. A compound according to Claim 16 which is 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
32. A compound according to Claim 16 which is 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
33. A compound according to Claim 16 which is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
34. A compound according to Claim 16 which is 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)imidazo[4,5-c]pyridine.
35. A compound according to Claim 16 which is 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine.
36. A compound according to Claim 16 which is 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine.
37. A compound according to Claim 16 which is 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
38. A compound according to Claim 16 which is 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
39. A compound according to Claim 16 which is 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
40. A compound according to Claim 16 which is 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
41. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine.
42. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine.
43. A compound according to Claim 16 which is 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
44. A compound according to Claim 16 which is 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
45. A compound according to Claim 1 wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring or phenyl. Y is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms. n is an integer of 1 to 3.  $R_3$  is hydrogen or alkyl of 1 to 6 carbon atoms and  $R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.
46. A compound according to Claim 45 wherein the halogen is fluoro.
47. A compound according to Claim 45 which is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.



48. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.

49. A compound according to Claim 45 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.

50. A compound according to Claim 45 wherein the alkoxy is methoxy.

51. A compound according to Claim 45 which is 5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.

52. A compound according to Claim 45 which is 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.

53. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.

54. A compound according to Claim 45 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.

55. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride.

56. Use of at least one compound according to Claim 1 for preparing a medicament for treating diseases or disorder mediated by platelet-activating factor.


57. Use according to Claim 56 wherein said compound is selected from the group consisting of

- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-n-octylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-n-decylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-n-dodecylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-2-decyl-N-methylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)imidazo[4,5-c]pyridine
- 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine
- 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine
- 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)-2-methylbenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride
- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine
- 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine
- 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine
- 5-[4-{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-isopropyl,N-3-methylcyclopentyl)carboxamido}benzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine
- 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine

5-[4-{-(N-cyclopentyl,N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-{-(N-isopropyl,N-cyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine

5 Amended claims in accordance with Rule 86(2) EPC.

1. A compound of the formula



or a pharmaceutically acceptable acid addition thereof; wherein

R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl or 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; heterocycloalkyl having 4 to 8 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; heteroaromatic having 5 or 6 carbon atoms which can be optionally substituted by alkyl of 1 to 6 carbon atoms; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen; R<sub>1</sub> and R<sub>2</sub> cannot both be hydrogen


Y is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen; thioalkyl wherein the alkyl is 1 to 6 carbon atoms; alkoxyalkyl wherein the alkyl groups are each 1 to 6 carbon atoms; hydroxyalkyl wherein the alkyl is 1 to 6 carbon atoms; alkylthioalkyl wherein the alkyl group are each 1 to 6 carbon atoms; cyano; mercaptoalkyl wherein the alkyl is 1 to 6 carbon atoms; hydroxy; amino; alkylamino wherein the alkyl group are each 1 to 6 carbon atoms; and dialkylamino wherein the alkyl group are each 1 to 6 carbon atoms.

n is an integer of 1 to 5.

R<sub>3</sub> is a group substituted at one or more of the 4, 6, or 7 positions of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms.

2. A compound according to Claim 1 having the formula



or a pharmaceutically acceptable acid addition salt thereof; wherein

R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15

carbon atoms; cycloalkyl having 3 to 8 carbon atoms; substituted cycloalkyl which can be substituted one or more by alkyl of 1 to 6 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; phenyl; substituted phenyl which can be substituted one or more by a group independently selected from alkyl of 1 to 6 carbon atoms or halogen; straight or branched alkenyl having 3 to 15 carbon atoms with the proviso that the double bond of the alkenyl group cannot be adjacent to the nitrogen; cycloalkenyl having 5 to 8 carbon atoms with the proviso that the double bond cannot be adjacent to the nitrogen;  $R_1$  and  $R_2$  cannot both be hydrogen

$Y$  is phenyl or phenyl substituted once or more than at one or more of the 2, 3, 5 or 6 position of the phenyl ring by substituents independently selected from the group consisting of alkoxy wherein the alkyl is 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro, or chloro; straight or branched chain alkyl having 1 to 6 carbon atoms; substituted straight or branched chain alkyl which can be substituted one or more by halogen;

$n$  is an integer of 1 to 5.

$R_3$  is a group substituted at one or more of the 4, 6, or 7 position of the pyridine ring said group being independently selected from hydrogen; alkyl of 1 to 6 carbon atoms; halogen wherein the halogen is selected from bromo, fluoro or chloro; alkoxy wherein the alkyl is 1 to 6 carbon atoms.

$R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

3. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)-carboxamido}benzyl]imidazo[4,5-c]pyridine.

4. A compound according to Claim 2 which is 5-[4{-(N-isopropyl,N-3-methylcyclopentyl)-carboxamido}benzyl]imidazo[4,5-c]pyridine.

5. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)-carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine.

6. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl,N-3-methylcyclohexyl)-carboxyamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.

7. A compound according to Claim 2 which is 5-[4{-(N-cyclopentyl-N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.

8. A compound according to Claim 2 which is 5-[4{-(N-isopropyl,N-cyclohexyl) carboxamido}benzyl]imidazo[4,5-c]pyridine.

9. A compound according to Claim 2 wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl.

10. A compound according to Claim 9 wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; or cycloalkyl having 3 to 8 carbon atoms.

11. A compound according to Claim 2 where  $Y$  is phenyl.


12. A compound according to Claim 2 wherein  $Y$  is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms.

13. A compound according to Claim 12 wherein the halogen is fluoro.

14. A compound according to Claim 12 wherein the alkoxy is methoxy.

15. A compound according to Claim 2 wherein  $n$  is an integer of 1 to 3.

16. A compound according to Claim 1 having the formula



or a pharmaceutically acceptable acid addition salt thereof: wherein  $R_1$  and  $R_2$  are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring; or phenyl.  $n$  is an integer of 1 to 3.  $R_3$  is hydrogen or alkyl of 1 to 6 carbon atoms and  $R_4$  is hydrogen or alkyl of 1 to 6 carbon atoms.

17. A compound according to Claim 16 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
18. A compound according to Claim 16 which is 5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
- 5 19. A compound according to Claim 16 which is 5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
20. A compound according to Claim 16 which is 5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
21. A compound according to Claim 16 which is 5-[4-(N-2-decalyl)-N-methylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
- 10 22. A compound according to Claim 16 which is 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
23. A compound according to Claim 16 which is 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
- 15 24. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
25. A compound according to Claim 16 which is 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
26. A compound according to Claim 16 which is 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
- 20 27. A compound according to Claim 16 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
28. A compound according to Claim 16 which is 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
- 25 29. A compound according to Claim 16 which is 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
30. A compound according to Claim 16 which is 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
31. A compound according to Claim 16 which is 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
- 30 32. A compound according to Claim 16 which is 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
33. A compound according to Claim 16 which is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
- 35 34. A compound according to Claim 16 which is 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)]imidazo[4,5-c]pyridine.
35. A compound according to Claim 16 which is 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine.
36. A compound according to Claim 16 which is 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine.
- 40 37. A compound according to Claim 16 which is 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
38. A compound according to Claim 16 which is 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
- 45 39. A compound according to Claim 16 which is 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
40. A compound according to Claim 16 which is 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine.
41. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine.
- 50 42. A compound according to Claim 16 which is 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine.
43. A compound according to Claim 16 which is 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]-imidazo[4,5-c]pyridine.
- 55 44. A compound according to Claim 16 which is 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine.

45. A compound according to Claim 1 wherein R<sub>1</sub> and R<sub>2</sub> are each independently selected from hydrogen; straight or branched chain alkyl of 1 to 15 carbon atoms; cycloalkyl having 3 to 8 carbon atoms; bicycloalkyl having 3 to 8 carbon atoms in each ring or phenyl. Y is substituted phenyl wherein the substituent is halogen selected from the group consisting of bromo, fluoro, or chloro or alkoxy wherein the alkyl is 1 to 6 carbon atoms. n is an integer of 1 to 3. R<sub>3</sub> is hydrogen or alkyl of 1 to 6 carbon atoms and R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms.
46. A compound according to Claim 45 wherein the halogen is fluoro.
47. A compound according to Claim 45 which is 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
48. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
49. A compound according to Claim 45 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine.
50. A compound according to Claim 45 wherein the alkoxy is methoxy.
51. A compound according to Claim 45 which is 5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
52. A compound according to Claim 45 which is 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
53. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
54. A compound according to Claim 45 which is 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine.
55. A compound according to Claim 45 which is 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride.
56. A pharmaceutical composition useful for treating diseases or disorder mediated by platelet-activating factor comprising at least one compound according to Claim 1, together with one or more non-toxic pharmaceutically acceptable carriers.
57. A pharmaceutical composition according to Claim 56 wherein said compound is selected from the group consisting of
- 5-[4-(N-methyl-N-cyclohexylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-n-octylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-n-decylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-n-dodecylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-2-decyl-N-methylcarboxamido)-benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclohexylcarboxamidobenzyl)imidazo[4,5-c]pyridine
  - 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine
  - 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-phenyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-3-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-4-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N,N-dicyclopentylcarboxamido)-2-methylbenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine
  - 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride
  - 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine

- 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine  
 5-[4-{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine  
 5-[4-{-(N-isopropyl,N-3-methylcyclopentyl)carboxamido}benzyl]imidazo[4,5-c]pyridine  
 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-{-(N-cyclopentyl,N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-{-(N-isopropyl,N-cyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine  
 58. Use of a compound of Claim 1 for preparing a medicament for treating diseases or disorder  
 mediated by platelet-activating factor in a mammal in need of such treatment.  
 59. Use according to Claim 58 wherein said compound is selected from the group consisting of  
 5-[4-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-n-octylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-n-decylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-n-dodecylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-2-decyl-N-methylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-2-(2,4,4-trimethyl)pentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-diisopropylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-ethyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-sec.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-3-pentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclobutyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclohexylcarboxamido)imidazo[4,5-c]pyridine  
 5-[2-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]ethyl]imidazo[4,5-c]pyridine  
 5-[3-[4-(N-methyl-N-cyclohexylcarboxamido)phenyl]propyl]imidazo[4,5-c]pyridine  
 5-[4-(N-tert.butyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-phenyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-3-N-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-4-N-methylcyclohexyl-N-cyclopentylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-cyclohexyl-N-cyclopentylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine, hydrochloride  
 5-[4-(N-cyclopentyl-N-cyclohexylcarboxamido)-2-methoxybenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-isopropyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[4-(N-methyl-N-cyclohexylcarboxamido)-2-fluorobenzyl]imidazo[4,5-c]pyridine  
 5-[3-(N-methyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[3-(N-isopropyl-N-cyclohexylcarboxamido)benzyl]imidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-4-methylimidazo[4,5-c]pyridine  
 5-[4-(N,N-dicyclopentylcarboxamido)benzyl]-2-methylimidazo[4,5-c]pyridine.  
 5-[4-{-(N-cyclopentyl,N-3,5-dimethylcyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine.  
 5-[4-{-(N-isopropyl,N-3-methylcyclopentyl)carboxamido}benzyl]imidazo[4,5-c]pyridine.  
 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-3-methoxybenzyl]imidazo[4,5-c]pyridine.  
 5-[4-{-(N-cyclopentyl,N-3-methylcyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.  
 5-[4-{-(N-cyclopentyl,N-cyclohexyl)carboxamido}-2-methoxybenzyl]imidazo[4,5-c]pyridine.  
 5-[4-{-(N-isopropyl,N-cyclohexyl)carboxamido}benzyl]imidazo[4,5-c]pyridine.

60. A method for treating diseases or disorder mediated by platelet-activating factor comprising administering a therapeutically effective dose of pharmaceutical composition of Claim 56 to mammal in need of such treatment.

5

10

15

20

25

30

35

40

45

50

55



European Patent  
Office

**PARTIAL EUROPEAN SEARCH REPORT**  
which under Rule 45 of the European Patent Convention  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

Application number

EP 89104464.6

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
P, A	EP - A - 0 260.613 (G.D. SEARLE & CO) * examples 3-5, 11, 15; page 21, table I; abstract *; & US - A - 4804858 (Cat. D) ---	1, 2, 16, 56	C07D471/04 A61K31/435 //(C07D471/04 235:00, 221:00)
A	EP - A - 0093593 (E. LILLY AND CO) * abstract * ---	1, 56	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
			C07D471/00
<b>INCOMPLETE SEARCH</b>			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-57 Claims searched incompletely: 58-60: Methods for treatment of the human or animal body by surgery or therapy (Art. 52(4) EPC) Claims not searched: Reason for the limitation of the search:</p>			
Place of search Berlin		Date of completion of the search 06.06.89	Examiner HASS, C.V.F.
<b>CATEGORY OF CITED DOCUMENTS</b>			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO Form 1505.1 (01/82)